

Elena Marco-Herrero¹, J. Ignacio González-Gordillo² and José A. Cuesta¹

Morphology of the megalopa of the mud crab, *Rhithropanopeus harrisii* (Gould, 1841) (Decapoda, Brachyura, Panopeidae), identified by DNA barcode

¹ Instituto de Ciencias Marinas de Andalucía, CSIC. Avda. República Saharaui, 2, 11519 Puerto Real, Cádiz, Spain. elena.marco@csic.es, jose.cuesta@icman.csic.es

² Campus de Excelencia Internacional del Mar (Ceimar), Departamento de Biología, Universidad de Cádiz, 11510 Puerto Real, Cádiz, Spain. nacho.gonzalez@uca.es

Corresponding author: elena.marco@csic.es, Tlf: +34.956 83 26 12 (ext. 324) Fax: +34. 956 83 47 01

Abstract

The morphology of the megalopa stage of the panopeid *Rhithropanopeus harrisii* is redescribed and illustrated in detail from plankton specimens identified by DNA barcode (16S mtDNA) as previous descriptions do not meet the current standard of brachyuran larval description. Several morphological characters vary widely from those of other panopeid species which could cast some doubt on the species' placement in the same family. Besides, some anomalous megalopae of *R. harrisii* were found among specimens reared at the laboratory from zoeae collected in the plankton. These anomalous morphological features are discussed in terms of problems associated with laboratory rearing conditions.

Keywords *Rhithropanopeus harrisii*, Panopeidae, Megalopa, Barcode, 16S, Morphology, Anomalies.

Introduction

Currently three species of Panopeidae are known for the Iberian Peninsula, *Panopeus africanus* A. Milne-Edwards, 1867, *Dyspanopeus sayi* (Smith, 1869) and *Rhithropanopeus harrisii* (Gould, 1841). While *Panopeus africanus* is an Iberian native species distributed from the Gulf of Cadiz (SW Spain) to the Mondego estuary (NW Portugal), the other two panopeids are introduced species. These are among the most widespread introduced brachyuran species in the world. *Dyspanopeus sayi* is native to the Atlantic coast of North America from Florida to Canada (Nizinski 2003), and has

been introduced to coastal areas of south-west England, Queens Dock, Swansea (Wales) (Ingle 1980;Clark 1986), to the French and Dutch coasts of the North Sea (Vaz et al. 2007), the Black Sea (Micu et al. 2010), and more recently to the Mediterranean Sea, Venice, the Marano and Varano lagoons, the Po River Delta (western Adriatic Sea) (Frogliia and Speranza 1993;Mizzan 1995; Florio et al.2008) and to the east coast of the Iberian Peninsula (Schubart et al. 2012).The first report of a population of *R. harrisii* for the Iberian Peninsula was made by Cuesta et al. (1991) for the Guadalquivir estuary, but populations are present in many European Atlantic estuaries, as well as in some Mediterranean locations. The species has been extensively studied from several perspectives such as ecology, phylogeography and larval biology (Gonçalves et al. 1995; Forward 2009; Projecto-García et al. 2010).

Rhithropanopeus harrisii is a euryhaline crab typically associated with sheltered estuarine habitats. Connolly (1925) described its four zoeal stages and the megalopa, based on larvae reared from eggs in the laboratory. Further descriptions were provided by Hood (1962) and Chamberlain (1962), but the best illustrations of the larval stages are shown in Costlow and Bookhout (1971) (as underlined by Forward 2009). Nevertheless, all descriptions are incomplete compared to the current standard of brachyuran larval descriptions proposed by Clark et al. (1998).

Traditionally, descriptions of larvae have been accomplished from specimens cultivated in the laboratory under controlled conditions (temperature, salinity, density and absence of predators), and the specimens commonly originated from a single or sometimes from two ovigerous females. These circumstances may contribute to conceal the morphological variability of larvae that can be found in the field, a phenomenon already discussed in the literature for brachyuran larvae (Cuesta et al. 2002).

The use of molecular markers has demonstrated to be a powerful tool in providing accurate identifications for plankton specimens (Pan et al. 2008; Pardo et al. 2009; Ampuero et al. 2010; Marco-Herrero et al. 2013a). The identification of megalopae has traditionally been based on morphological characteristics, but sometimes it is impossible to get an accurate identification with this approach. In the present study we used partial sequences of the mitochondrial gene 16S as DNA barcode to identify the megalopae collected in the plankton. The 16S marker has proven to be an effective tool in studies of decapod crustaceans (Schubart et al. 2000; Porter et al. 2005; Ahyong et al. 2007), not only for the establishment of new species, but also to elucidate the taxonomic validity of closely related species (Schubart et al. 1998, 2001; Spivak and Schubart 2003).

In contrast to traditional descriptions, the megalopae of the present study were obtained from the plankton and identified by DNA barcode. Furthermore, in order to provide a definite morphological description of the megalopa stage of *R. harrisii*, comparisons were made not only with previous descriptions, but also with another set of megalopae which were reared in the laboratory from four zoeae I collected in the plankton.

Materials and Methods

Collection of the megalopae

Twenty-eight megalopae of *R. harrisii* were collected in July 2007 and four zoeae I in April 2011, all from the plankton of the Guadalete estuary (Cádiz-SW Spain) (36°35'24.09"N 6°13'46.19"W).

Rearing and description of the megalopae

All megalopae collected were preserved directly in 80% ethanol. The four zoeae I were placed in beakers containing filtered and well-aerated sea water at a salinity of 32 ± 1 ‰ and a temperature of 26 ± 1 °C. The larvae were fed with the rotifer *Brachionus plicatilis* (fed with *Nannochloropsis gaditana*). Rearing was finished when all zoeae had moulted to the megalopa instar. Megalopa descriptions were based on 10 specimens identified by DNA barcode.

To facilitate the microscopical observation of larvae structures, a digestion-stain procedure was carried out. Firstly, entire specimens were placed for about 10 minutes in a watch glass with 2 ml of heated lactic acid. Immediately afterwards, three drops of Clorazol Black stain (0.4 g Clorazol Black powder dissolved in 75 ml 70% EtOH) were added to the heated solution. After 5-10 minutes, the specimen was removed from the solution and placed on a slide with lactic acid in order to proceed with the dissection of the appendages (Landeira et al. 2009).

Drawings and measurements were made using a Leica MZ6 and Zeiss Axioskop compound microscope with Nomarski interference, both equipped with a *camera lucida*. All measurements were made by an ocular micrometer. The measurements taken were: cephalothorax length (CL) as the distance from the tip of the rostrum to the posterior margin of the cephalothorax; cephalothorax width (CW) as the maximum

width of the carapace. Two megalopae identified by DNA barcode were deposited at the Biological Collections of Reference of the Institut de Ciències del Mar (ICM-CSIC) in Barcelona, under accession numbers ICMD13121701 and ICMD13121702.

DNA extraction, amplification and sequencing

The identification of the megalopae was based on partial sequences of the 16S rDNA gene. Total genomic DNA was extracted from muscle tissue from 1-2 pereopods of each megalopa, and incubated for 1–24 hours in 300 µl lysis buffer at 65°C. Protein was precipitated by addition of 100 µl of 7.5M ammonium acetate and subsequent centrifugation, and DNA precipitation was obtained by addition of 300 µl isopropanol and posterior centrifugation. The resulting pellet was washed with ethanol (70%), dried, and finally resuspended in Milli-Q distilled water.

Target mitochondrial DNA from the large subunit rRNA (16S) gene was amplified with polymerase chain reaction (PCR) and the following cycling conditions for reactions: 2 min at 95°C, 40 cycles of 20 s at 95°C, 20 s at 45-48°C, 45 s at 72°C, and 5 min at 72°C. Primers 1472 (5'-AGA TAG AAA CCA ACC TGG -3') (Crandall and Fitzpatrick 1996) and 16L2 (5'-TGC CTG TTT ATC AAA AAC AT-3') (Schubart et al. 2002) were used to amplify 540 bp of 16S. PCR products were sent to NewBiotechnic and Biomedal companies to be purified and then two-direction sequencing.

Sequences were edited using the software Chromas version 2.0. The final sequences were blasted on Genbank database to get the best Blast matches for an accurate identification. Sequences are accessible in Genbank under the accession numbers KJ125076-KJ125077.

Results

Barcode identification

Using the BLAST utility (<http://blast.ncbi.nlm.nih.gov/Blast.cgi>), the sequences obtained from the megalopae were compared with those deposited in GenBank. The sequences perfectly fit those of *Rhithropanopeus harrisii*, more specifically, no difference (100% match) was found between the 16S sequence for 546 bp and sequences of *R. harrisii* from Woodland Beach, Delaware, USA (U11Z 3836), Genbank accession number AJ274697.

Nevertheless, three out of four megalopae reared in the laboratory from specimens collected as zoeae I in the plankton, did not show the general morphology and all setation patterns of those megalopal stage of *Rhithropanopeus harrisii* which had been directly collected in the plankton. According to the DNA barcode, however, these specimens clearly belong to the same species. We have considered these specimens as ‘anomalous megalopa’ and have provided an additional description of this type of larva.

Description of the megalopa

(Figs 1A-E; 2A, D, E, G; 3A, C, D; 4A-E; 5A, D, E)

Size: CL = 1.18 ± 0.05 mm; CW = 1.02 ± 0.05 mm; N= 5

Cephalothorax (Figs 1A, B): Rostrum is short and obliquely downward with 2 lateral simple setae at base, anterior end with a median triangular notch; the pedunculated

eyes with 8 small simple setae each; hepatic region swollen; one pair each of protogastric, mesobranchial and cardiac protuberances present; broader posterior part, margins setose.

Antennule (Fig. 2A): Peduncle three-segmented, with 3 short simple setae on first segment, 2 short simple setae on median segment and 2 short simple setae plus 2 pairs of long plumodenticulate setae on distal segment; endopod unsegmented with 1 basal simple seta, 1 subterminal simple seta and 3 terminal simple setae; exopod three-segmented, with 10 aesthetascs (arranged 0, 4, 6) and 4 setae (arranged 0, 2, 2 setae).

Antenna (Fig. 2B): Peduncle three-segmented with 6 setae (arranged 4, 1, 1); flagellum six-segmented with 10 simple setae (arranged 0, 0, 1, 4, 3, 2).

Mandible (Fig. 2D): Palp two-segmented, with 5 terminal short plumodenticulate setae on distal segment.

Maxillule (Fig. 2E): Coxal endite with 12 plumose setae; basal endite with 16 setae (3 terminal plumodenticulate, 1 terminal sparsely plumose, 7 terminal cuspidate, 3 subterminal plumodenticulate, and 2 proximal plumodenticulate); endopod unsegmented with 1 proximal and 2 terminal simple setae; long epipodal seta present.

Maxilla (Fig. 2G): Coxal endite bilobed with 2 + 3 terminal plumose setae; basal endite bilobed with 6 + 6 sparsely plumodenticulate setae; endopod unsegmented and without setae; scaphognathite with 45-47 marginal plumose setae plus 2 small simple setae on each lateral surface.

First maxilliped (Fig. 3A): Epipod well developed, triangular shaped, with 5 long simple setae and 1 proximal plumodenticulate seta; coxal endite with 5 inner simple

setae and 7 terminal plumose setae; basal endite with 1 inner + 4 subterminal + 11 terminal sparsely plumodenticulate setae plus 2 terminal short simple setae; endopod unsegmented with 4 short terminal simple setae; exopod two-segmented, with 5 long terminal plumose setae on distal segment.

Second maxilliped (Fig. 3C): Reduced epipod with 2 simple setae and 1 plumodenticulate seta; endopod five-segmented, with 1 simple, 2 simple, 1 simple, 4 plumodenticulate + 1 short simple, and 3 proximal simple + 6 terminal plumodenticulate setae, respectively; exopod two-segmented, with 2 simple setae on proximal segment and 5 long terminal plumose setae on distal one.

Third maxilliped (Fig. 3D): Epipod well developed with a proximal marginal row of 6 plumose setae and 14 long simple setae; protopod with a marginal row of 7 plumose setae and 1 simple + 3 plumose inner setae; endopod five-segmented, with 19, 14, 6, 9, and 9 setae respectively; exopod two-segmented with 5 long plumose setae on distal segment.

Pereiopods (Figs 4A-E): Pereiopods 2-5 thin and setose, with long subterminal setae on dactyli. Cheliped robust and setose without remarkable recurved spines, only sometimes a small spine, never recurved.

Sternum (Figs 1D, E): Maxilliped sternites completely fused with 6 simple setae, cheliped sternites with 4 or 6 simple setae each, pereopod sternites 2-5 with 3 or 4, 2 or 3, 1 or 2, and 0 simple setae, respectively; sternal sutures are interrupted medially. There are two forms according to setation; the most common is illustrated in Fig. 1.

Pleon (Fig. 5A): Six somites plus telson; setation as show.

Pleopods (Figs 5D, E): Biramous except uropods, present on somites 2-5; endopod with 3 cincinnuli; exopod with 10 long plumose natatory setae; uropod with 3 or 4 natatory setae on distal segment.

Description of anomalous megalopae

(Figs 1F, 2C, F, H, 3B, 5B, C)

Size: CL = 1.12-1.14 mm; CW = 0.92-0.98 mm; N= 2

All three specimens exhibited the following deviations from the typical form: cephalothorax with different shape, bearing vestiges of zoeal lateral spines, and a reduced number of setae (Fig. 1F); antennular peduncle with remains of exopodal and protopodal processes as spines (Fig. 2C); endopod of maxillule with a setation pattern of 1,2,2,2 as in the zoeal endopod of the maxillule (Fig. 2F); endopod of maxilla with setation 3,2,2 as in the zoeal maxillar endopod (Fig. 2H); endopod of first maxilliped with 3 terminal long setae plus 1+1+1 long inner plumose setae (Fig. 3B); telson with 2-3 terminal setae in the place of furcal arms and 1 pair of marginal setae as zoeal stage (Figs 5B, C).

Discussion

Redescriptions of brachyuran larval stages are unusual, although they are necessary when previous descriptions are brief, incomplete, inaccurate or deficient, making them useless for reliable identifications. There are some cases of redescriptions

in the recent literature. For instance, *Aratus pisonii* (H. Milne Edwards, 1837) was redescribed by Cuesta et al. (2006) considering that the previous description by Warner (1968) referred to a clearly anomalous megalopa. The most recent redescription of *Dyspanopeus sayi* by Marco-Herrero et al. (2013b) was necessary because the several previous descriptions were brief and inaccurate and thus inappropriate for comparative taxonomic studies. Correct descriptions of larval stages are needed for phylogenetic studies and accurate identifications of plankton-collected specimens. In the case of *R. harrisii*, the several previous descriptions of the megalopa from both laboratory reared larval stages and from plankton-collected specimens are all incomplete and inaccurate and do not meet the standard proposed by Clark et al. (1998), currently followed by the majority of decapod larval morphologists.

Since the previous descriptions do not allow for an accurate identification of plankton-collected specimens, the DNA barcode was used instead. Current molecular tools ensure a correct identification of specimens collected in the field, which present clear advantages over specimens which have been reared in the laboratory. In particular, field-collected larvae allow for obtaining a better representation of natural morphological variability compared to larvae originated from only one or two ovigerous females cultured in the laboratory. In the present study the 16S sequences of the 10 studied megalopae, collected in the Guadalete estuary for morphological description, fit at 100% the 16S sequence of *R. harrisii* from Delaware (USA) deposited in Genbank.

The morphology of the megalopae of *R. harrisii* described in the present work do not completely match the typical characters of the megalopa stages of panopeids, although Martin (1984), based on zoeal morphology, included *R. harrisii* in the Group I together with the majority of panopeids. Even when the classification was based on megalopal features, the species was attributed to Group I (Martin 1988). The main

differences relate to rostrum morphology, the number of segments of the antennular flagellum, and to the spinulation of the ischium of the cheliped.

The typical panopeid megalopa rostrum presents a remarkable spine at each basal angle, called “horns” in some papers, but these are missing in *R. harrisii*. The antennular flagellum of *R. harrisii* shows six segments while eight segments are present in other panopeids such as *D. sayi* (see Marco-Herrero et al. 2013b) and *P. africanus* (see Rodriguez and Paula1993). The number of segments of the antennular flagellum is considered to be a conservative character at family level in other taxa (Cuesta 1999). Finally, the absence of a remarkable recurved spine on the cheliped ischium is another marked contrast to the majority of panopeids. Together with the above-mentioned differences, this feature could challenge the phylogenetic position of this species. Future molecular phylogenetic studies will help to resolve this question raised by the larval morphology.

The setation patterns of maxillule, maxilla, first, second and third maxillipeds, and sternum are described in the present work for the first time. As to the setation pattern of the sternal plates some variability was observed, although the proportions between sternites were always similar.

In the identification key to the megalopa stages of the Mediterranean Brachyura by Pessani et al. (2004), *R. harrisii* is differentiated by bearing three long plumose terminal setae on the distal segment of the uropod in contrast to “uropod exopod with more than 3 setae”. Megalopae in the present study showed either three or four setae, and in one case this variability occurred in the same specimen. The same variability in the setation on the exopods of the uropods has already been described by Kurata (1970).

In the present work we also studied megalopae grown from zoeae I which had been collected in the plankton and raised in the laboratory. There is some evidence that the culture conditions (temperature and/or feeding) were suboptimal. The megalopae which developed under these conditions showed an anomalous morphology. This kind of anomalies has already been reported in other species, and not only for larvae raised in the laboratory (Willems 1982; Cuesta and Anger 2001), but also for larvae collected in the field (Cuesta et al. 2002). In all these cases, the anomalies referred to morphological character of the zoeal phase, such as the presence of short lateral spines in the cephalothorax and the setation patterns of maxillule and maxilla endopods. The available data suggest that morphological anomalies in the megalopa stage are the result of suboptimal environmental conditions (temperature, salinity, food), and that such deficiencies can occur not only during laboratory rearing but also in the natural environment.

Acknowledgements

We want to thank the “Real Club Náutico” of the Puerto Santa María (Cádiz, Spain) for kindly allowing us to use their facilities. We are especially grateful to Reyes Sánchez García and Ainhoa Olaguer-Feliú who helped in collecting the larvae, Carlos Sánchez Nieto for his assistance in the laboratory work, and Dulce Subida for her careful language corrections. Thanks are also due to anonymous referees and Prof. Heinz-Dieter Franke for his editorial work. The research was carried out within the framework of the projects MEGALOPADN (CGL2009-11225) and “Transporte y reclutamiento larvario de crustáceos bentónicos litorales: importancia de los agentes forzadores

costeros y regimen mareal” (CTM2005-00024/MAR), both co-financed by the “Ministerio de Economía y Competividad (MINECO)” Spanish Plan R+D+I and European Union FEDER funds. Elena Marco-Herrero acknowledges the pre-doctoral FPI Fellowship supported by MINECO (BES-2010-033297).

References

Ampuero D, Palma AT, Veliz D, Pardo LM (2010) Description, seasonal morphological variation, and molecular identification of *Paraxanthus barbiger* megalopae obtained from the natural environment. *Helgol Mar Res* 64:117–123

Ahyong ST, Lai JCY, Sharkey D, Colgan DJ, Ng PKL (2007) Phylogenetics of the brachyuran crabs (Crustacea: Decapoda): The status of Podotremata based on small subunit nuclear ribosomal RNA. *Mol Phylogenet Evol* 45:576–586

Chamberlain NA (1962) Ecological studies on the larval development of *Rhithropanopeus harrisii* (Xanthidae, Brachyura). Chesapeake Bay Institute, Johns Hopkins Univ Tech Rep 28:1–47

Clark PF (1986) North-east Atlantic crabs: an atlas of distribution. Marine Conservation Society. England

Clark PF, Calazans DK, Pohle GW (1998) Accuracy and standardization of brachyuran larval descriptions. *Invert Reprod Dev* 33:127–144

Connolly CJ (1925) The larval stages and megalops of *Rhithropanopeus harrisii* (Gould). *Contrib Can Biol Stud Biol Stn Can* 2:329–334

321 Costlow JD, Bookhout CG (1971) The effect of cyclic temperatures on larval
 322 development in the mud-crab *Rhithropanopeus harrisi*. In: Crisp DJ (ed) Fourth
 323 European Marine Biology Symposium. Cambridge University Press, Cambridge, pp.
 324 211–220

325 Crandall KA, Fitzpatrick JFJ (1996) Crayfish molecular systematics: Using a
 326 combination of procedures to estimate phylogeny. *SystBiol* 45:1–26

327 Cuesta JA (1999) Morfología larval de la familia Grapsidae (Crustacea, Decapoda,
 328 Brachyura). Dissertation, University of Seville, Spain

329 Cuesta JA, García-Raso JE, González-Gordillo JI (1991) Primera cita de
 330 *Rhithropanopeus harrisi* (Gould, 1814) (Crustacea, Decapoda, Brachyura, Xanthidae)
 331 en la Península Ibérica. *Bol Inst Esp Oceanogr* 7:149–153

332 Cuesta JA, Anger K (2001) Larval morphology of the sesarmid
 333 crab *Armases angustipes* (Dana, 1852) (Crustacea, Decapoda, Grapsoidea) reared under
 334 laboratory conditions. *J Crust Biol* 21:821–838

335 Cuesta JA, Luppi TA, Rodríguez A, Spivak ED (2002) Morphology of the megalopal
 336 stage of *Chasmagnathus granulatus* Dana, 1851 (Crustacea: Decapoda: Brachyura:
 337 Varunidae), with comments on morphological anomalies. *Proc Biol Soc Wash* 115:391–
 338 402

339 Cuesta JA, García-Guerrero MU, Rodríguez A, Hendrickx ME (2006) Larval
 340 morphology of the sesarmid crab *Aratus pisonii* (H. Milne Edwards, 1837) (Decapoda,
 341 Brachyura, Grapsoidea) from laboratory reared material. *Crustaceana* 79:175–196

342 Florio M, Breber P, Scirocco T, Specchiulli A, Cilenti L, Lumare L (2008) Exotic
 343 species in Lesina and Varano lakes, Gargano National Park (Italy). *Transit Waters Bull*
 344 2:69–79

345 Forward RB Jr (2009) Larval biology of the Crab *Rhithropanopeus harrisii* (Gould): A
 346 synthesis. *Biol Bull* 216:243–256

347 Froggia C, Speranza S (1993) First record of *Dyspanopeus sayi* (Smith, 1869) in the
 348 Mediterranean Sea (Crustacea: Decapoda: Xanthidae). *Quad Ist Ricerca Pesca Maritt*
 349 5:163–166

350 Gonçalves F, Ribeiro R, Soares AM (1995). *Rhithropanopeus harrisii* (Gould), an
 351 American crab in the estuary of the Mondego River, Portugal. *J Crust Biol* 15:756–762

352 Hood RM (1962) Studies on the larval development of *Rhithropanopeus harrisii*
 353 (Gould) of the family Xanthidae (Brachyura). *Gulf Res Rep* 1:122–130

354 Ingle R W (1980) *British Crabs*. Oxford University Press, London

355 Kurata H (1970) Studies on the life histories of decapod Crustacea of Georgia; Part III.
 356 Larvae of decapod Crustacea of Georgia. Dissertation, University of Georgia

357 Landeira JM, Lozano-Soldevilla F, González-Gordillo JI (2009) Morphology of first
 358 seven larval stages of the striped soldier shrimp, *Plesionika edwardsii* (Brandt, 1851)
 359 (Crustacea: Decapoda: Pandalidae) from laboratory reared material. *Zootaxa* 1986:51–66

360 Marco-Herrero E, Torres AP, Cuesta JA, Guerao G, Palero F, Abelló P (2013a) The
 361 systematic position of *Ergasticus* (Decapoda, Brachyura) and allied genera, a molecular
 362 and morphological approach. *Zool Scripta* 42:427–439

363 Marco-Herrero E, Guerao G, Cuesta JA (2013b) Morphology of the larval stages of a
 364 Mediterranean population of the allochthonous Say's mud crab, *Dyspanopeus sayi*
 365 (Decapoda: Brachyura: Panopeidae). Sci Mar 77:341–352

366 Martin JW, Felder DL, Truesdale FM (1984) A comparative study of morphology and
 367 ontogeny in juvenile stages of four western Atlantic xanthoid crabs (Crustacea:
 368 Decapoda: Brachyura). Philos Trans R Soc Lond B 303:537–604

369 Martin JW (1988) Phylogenetic significance of the brachyuran megalopa: Evidence
 370 from the xanthidae. Symp Zool Soc Lond 59:69–102

371 Micu D, Nita V, Todorova V (2010) First record of Say's mud crab *Dyspanopeus sayi*
 372 (Brachyura: Xanthoidea: Panopeidae) from the Black Sea. Mar Biodiver Rec 3:1–6

373 Mizzan L (1995) Notes on the presence and diffusion of *Dyspanopeus sayi* (Smith, 1869)
 374 (Crustacea, Decapoda, Xanthidae) in the Venetian Lagoon. Boll Mus Civ Stor
 375 Nat Venezia 44:121–130

376 Nizinski, MS (2003) Annotated checklist of decapod crustaceans of Atlantic coastal and
 377 continental shelf waters of the United States. Proc Biol Soc Wash 116:96–157

378 Pan M, McBeath A, Hay S, Pierce G, Cunningham C (2008) Real-time PCR assay for
 379 detection and relative quantification of *Liocarcinus depurator* larvae from plankton
 380 samples. Mar Biol 153:859–870

381 Pardo LM, Ampuero D, Veliz D (2009) Using morphological and molecular tools to
 382 identify megalopae larvae collected in the field: the case of sympatric *Cancer* crabs. J
 383 Mar Biol Ass UK 89:481–490

384 Pessani D, Tirelli T, Flagellal S (2004) Key for the identification of Mediterranean
 385 brachyuran megalopae. *Mediterranean Mar Sci* 5:53–64

386 Porter ML, Pérez-Losada M, Crandall K A (2005) Model-based multi-locus estimation
 387 of decapod phylogeny and divergence times. *MolPhylogenetEvol* 37:355–369

388 Projecto-Garcia J, Cabral H, Schubart CD (2010) High regional differentiation in a
 389 North American crab species throughout its native range and invaded European waters:
 390 a phylogeographic analysis. *Biol Invasions* 12:253–263

391 Rodríguez A, Paula J (1993) The larval and postlarval development of the mud crab
 392 *Panopeus africanus* A. Milne Edwards (Crustacea: Decapoda: Xanthidae), reared in the
 393 laboratory. *J Crust Biol* 13:296–308

394 Schubart CD, Diesel R, Hedges SB (1998) Rapid evolution to terrestrial life in Jamaican
 395 crabs. *Nature* 393:363–365

396 Schubart CD, Cuesta JA, Diesel R, Felder DL (2000) Molecular phylogeny, taxonomy,
 397 and evolution of nonmarine lineages within the Americangrapsoid crabs (Crustacea:
 398 Brachyura). *MolPhylogenetEvol* 15:179–190

399 Schubart CD, Cuesta JA, Rodríguez A (2001) Molecular phylogeny of the crab genus
 400 *Brachynotus* (Brachyura: Varunidae) based on the 16S rRNA gene.
 401 *Hydrobiology* 449:41–46

402 Schubart CD, Cuesta JA, Felder DL (2002) Glyptograpsidae, a new brachyuran family
 403 from Central America: Larval and adult morphology, and a molecular phylogeny of the
 404 Grapsoidea. *J Crust Biol* 22:28–44

405 Schubart CD, Guerao G, Abelló P (2012) First record and evidence of an established
 406 population of the North American mud crab *Dyspanopeus sayi*(Brachyura:
 407 Heterotremata: Panopeidae) in the western Mediterranean. Sci Mar 76:79–85

 408 Spivak ED,Schubart CD (2003) Species status in question: a morphometric and
 409 molecular comparison of *Cyrtograpsus affinis* and *C. altimanus* (Decapoda, Brachyura,
 410 Varunidae). J Crust Biol23:212–222

 411 Vaz S, Hofstede R, Martin J, Dewarumez JM, Verin Y, Le Roy D, Heessen, Daan N
 412 (2007) Benthic invertebrates community structure inferred from bottom trawl hauls
 413 observations and its relationships to abiotic conditions in the southern North Sea. ICES
 414 CM 2007/A :03

 415 Warner GF (1968) The larval development of the mangrove tree crab *Aratuspisonii*
 416 (H.Milne Edwards), reared in laboratory (Brachyura, Grapsidae). Crustaceana 2:249–
 417 258

 418 Willems KA (1982) Larval development of the land crab *Gecarcinuslateralislateralis*
 419 (Fréminville, 1835)(Brachyura: Gecarcinidae) reared in the laboratory.J Crust
 420 Biol2:180–201

 421

Figures captions

Fig.1 *Rhithropanopeus harrisii* (Gould, 1841).Megalopa, (A) frontal view; (B) dorsal view; (C) lateral view of the cephalothorax; (D,E) sternum; (F) anomalous megalopa, dorsal view

Fig.2*Rhithropanopeus harrisii* (Gould, 1841).Megalopa, (A) antennule; (B) antenna; (C) anomalous antenna;(D) mandible; (E)maxillule; (F)endopod of maxillule of the anomalous specimen; (G) maxilla; (H)endopod of maxillaof the anomalous specimen

Fig. 3 *Rhithropanopeus harrisii* (Gould, 1841).Megalopa, (A) first maxilliped; (B)endopod of first maxilliped of the anomalous specimen; (C) second maxilliped;(D) third maxilliped

Fig.4 *Rhithropanopeus harrisii* (Gould, 1841).Megalopa, (A) cheliped, with detail of the ischiumspine;(B) secondpereiopod; (C) thirdpereiopod; (D) fourthpereiopod; (E) fifthpereiopod

Fig. 5 *Rhithropanopeus harrisii* (Gould, 1841).Megalopa, (A): pleon, dorsal view; (B- C) telson of an anomalous megalopa; (D) uropod; (E) third pleopod